

L3(3.3)-Quadratic Relations in Factored Form

Key Concepts:

- factored form of quadratic relation
- direction of opening from 'a'
- solving for zeroes
- using symmetry to find:
 - x-coordinate of vertex
 - axis of symmetry
- using substitution to find:
 - y-coordinate of vertex
 - y-intercept

Is $y = 2(x+1)(x-5)$ a quadratic relation?

Examine 1st and 2nd differences:

x	y	1 st	2 nd
-2	14		
-1	0	$0 - (14) = -14$	$-10 - (-14) = 4$
0	-10	$-10 - (0) = -10$	$-6 - (-10) = 4$
1	-16	$-16 - (-10) = -6$	$-2 - (-6) = 4$
2	-18	$-18 - (-16) = -2$	

Quadratic
2nd differences
constant

The equation of a quadratic relation may be written in several forms:

- several forms:
1. standard form: $y = ax^2 + bx + c$ ← y-intercept
 2. factored form: $y = a(x - s)(x - t)$ ← zeroes
↓ method
 3. vertex form: $y = a(x - h)^2 + k$ ← vertex

The factored form, $y = a(x - s)(x - t)$, is most useful for finding the zeroes, which are $x = s$ and $x = t$.

Consider the following...

Give two numbers that have a product of zero:

★ $0(10) = 0$ $(0)(0) = 0$
 $10(0) = 0$ $\rightarrow (5+5) = 0$

What do you notice? $(\text{any value}) \times 0 = 0$

★ only way to get a product of zero is $x \cdot 0$


Solve:

(a) $\frac{3x}{3} = \frac{0}{3}$
 $x = 0$

(b) $\frac{57y}{57} = \frac{0}{57}$
 $y = 0$

(c) $\frac{3xy}{3} = \frac{0}{3}$
 $xy = 0$

Depending upon the location of the vertex, and whether the parabola opens up or down, it may have 0, 1, or 2 distinct (unique) zeroes.



Zeroes occur where the y-coordinate of the parabola is equal to zero.

To find the zeroes algebraically, we **set $y = 0$** and solve for the x-values that make the equation true.

Ex.1 Determine the zero(es) of each

(a) $y = x(x - 10)$

$$0 = x(x - 10)$$

\swarrow or \searrow
 $x = 0$ or $x = 10$

Recall:

Zero multiplied by anything is zero.

If $(a)(b) = 0$ then
 $a = 0$ or $b = 0$ (or both are zero).

(b) $y = -2(x - 5)(3x - 1)$

$$0 = -2(x - 5)(3x - 1)$$

\swarrow or \searrow
 $x = 5$ or $3x - 1 = 0$
 $\frac{3x}{3} = \frac{1}{3}$
 $x = \frac{1}{3}$
2 zeroes

(c) $y = 2(x - 2)^2$

$$0 = 2(x - 2)(x - 2)$$

$x - 2 = 0$ or $x - 2 = 0$
 $x = 2$ or $x = 2$

The zeroes and symmetry can be used to find the vertex (h, k).

Axis of Symmetry
AOS

For the x-coordinate (h), find the midpoint of the zeroes:

$$\star \quad MP = \frac{x_1 + x_2}{2} = \frac{s + t}{2}$$

For the y-coordinate (k), substitute the midpoint into the equation and solve for y:

$$\star \quad y = a(x - s)(x - t)$$

$$y = a(MP - s)(MP - t)$$

Steps to find Vertex
① find zeroes
② find AOS
③ sub AOS into equation to solve k

Ex.2 Determine the vertex:

(a) $y = -2(x - 2)(x - 8)$

$$\textcircled{1} \quad 0 = -2(x - 2)(x - 8)$$

$$\begin{array}{cc} \swarrow & \searrow \\ x - 2 = 0 & x - 8 = 0 \\ x = 2 & x = 8 \end{array}$$

$$\textcircled{2} \quad AOS = \frac{2 + 8}{2}$$

$$x = 5$$

$$\textcircled{3} \quad \text{Sub } x = 5 \text{ into equation}$$

$$= -2(5 - 2)(5 - 8)$$

$$= -2(3)(-3) \quad \therefore \text{the vertex is } (5, 18)$$

$$= 18$$

Ex.3 A parabola has zeroes at -3 and 2, and a y-intercept of 18. Determine the equation.

$$y = a(x - s)(x - t)$$

$$y = a(x - (-3))(x - 2)$$

$$y = a(x + 3)(x - 2)$$

$$18 = a(0 + 3)(0 - 2)$$

$$18 = a(3)(-2)$$

$$18 = -6a$$

$$\frac{18}{-6} = \frac{-6a}{-6}$$

$$a = -3$$

$$\therefore \text{the equation is}$$

$$y = -3(x + 3)(x - 2)$$

p. 155-157 # 2, 3, 4ace, 5, 6ace, 7, 10